### Designated Class2 OTC medicine

Please be sure to read these instructions before usage, and store them somewhere where they can be read if necessary.

# 血府逐瘀顆粒

## Keppu-Chikuo-karyu

This product is based on the formulation of "血府逐瘀湯(Keppu-chikuo-to)" which is described in 《Irin Kaisaku》, written during the Qing Dynasty. Relieves following symptoms of people past middle age who tend to have high blood pressure: headache, dull headache, stiff shoulder, dizziness, palpitations.



# [Precautions]



- 1. The following persons should contact a physician, pharmacist, or registered salesperson for a consultation before administration.
- (1) Patients undergoing medical treatment from a physician.
- (2) Pregnant women or women suspected of being pregnant.
- (3) Persons who have gastrointestinal weakness.
- 2. If the following symptoms are observed after taking this drug, these may be adverse reactions, so immediately discontinue the use of this drug, and show this document to your physician, pharmacist, or registered salesperson for a consultation.

Affected site	Symptom	
Gastrointestinal system	Loss of appetite, Gastric irritation	
other	Abnormal bleeding	

- 2. The following symptoms may be observed after taking this drug. If these symptoms persist or worsen, discontinue the use of this drug, and show this document to your physician, pharmacist, or registered salesperson for a consultation.

  Diarrhea, soft stool
- 3. When symptoms do not improve even after taking the medicine for about 1 month, stop taking the preparation and consult a physician, pharmacist or registered salesperson, being sure to take this instruction leaflet with you.

## [Indications]

Relieves following symptoms of people past middle age who tend to have high blood pressure: headache, dull headache, stiff shoulder, dizziness, palpitations.

# [Dosage and administration]

Three times a day, take it after the meal in 2-3 hours by the water or hot water.

Age	Single Dose	Doses per day	
Adult	1 nackaga (2.5g)	0 4:	
(15 years old and over)	1 package (2.5g)	3 times	
Children	Please do not take it.		
(under 15 years old)			

# [Ingredients (1day 3packages)]

An adult (15 years old and over) daily dosage of 3packages (7.5g) contains the below ingredients.

[Keppu-Chikuo-Ganryo-extract powder 3.0g]

Keppu-Chikuo-to-powder 1.5g

Ingredients		Quantities
トウキ Japanese Angelica root		1.98g
センキュウ	Cnidium Rhizome	0.99g
ジオウ	Rehmannia Root	1.98g
トウニン	Peach Kernel	2.64g
コウカ	Safflower	1.98g
キジツ	Immature Orange	1.32g
シャクヤク	Paony Root	1.32g
サイコ	Bupleurum Root	0.66g
カンゾウ	Glycyrrhiza	0.66g
キキョウ	Platycodon Root	0.99g
ゴシツ	Achyranthes Root	1.98g

Ingredients		Quantities
トウキ Japanese Angelica root		0.18g
センキュウ	Cnidium Rhizome	0.09g
ジオウ	Rehmannia Root	0.18g
トウニン	Peach Kernel	0.24g
コウカ	Safflower	0.18g
キジツ	Immature Orange	0.12g
シャクヤク	Paony Root	0.12g
サイコ	Bupleurum Root	0.06g
カンゾウ	Glycyrrhiza	0.06g
キキョウ	Platycodon Root	0.09g
ゴシツ	Achyranthes Root	0.18g

Additives: lactose, magnesium stearate

## < Precautions relating to ingredient and amount>

This product uses a natural ingredients (crude drugs) and as such look/taste slightly differently or they may stick together.

## [Precautions for storage and handling]

- (1) Store in a cool, low humidity place away from direct sunlight.
- (2) Keep out of reach of children.
- (3) Do not transfer the medicine to a different container.(It may lead to misuse or alter the quality of the drug)
- (4) Do not use any expired product.

## ■OTC 医薬品の多言語製品情報提供時に関する免責事項

- ・本製品は、日本国内での販売及び使用を目的として、日本の医薬品、医療機器等の品質、有効性及び 安全性の確保等に関する法律に基づき承認された医薬品です。
- ・多言語製品情報は、当該製品の日本語の添付文書を翻訳したものであり、あくまで参考情報として提供 されるものです。その記載内容及び当該製品自体が日本国外の法規に適合していることを保証するもの ではありません。
- ・多言語製品情報は、提供者(又は当社)による仮訳であり、今後、予告なしに修正・変更される可能性があります。
- ・多言語製品情報の記載内容に起因して発生した問題については、提供者(又は当社)は一切の責任を 負いかねます。

#### ■Disclaimer on Multilingual OTC Product Information

- This product is a pharmaceutical product approved under a Japanese law, the Law for Ensuring the Quality, Efficacy and Safety of Drugs and Medical Devices, with a view to its sale and use in Japan.
- Multilingual product information is a translation of the product labeling written in Japanese and provided for your information only. It does not warrant that its contents and the product itself conforms to laws and regulations in countries other than Japan.
- Multilingual product information is a tentative translation by the provider (or Our Company), and may be modified or altered without notice.
- The provider (or Our Company) assumes no responsibility for any occurred problem attributable to the contents of the multilingual product information.

Multilingual product information is a tentative translation by the provider (or Our Company), and may be modified or altered without notice.

The provider (or Our Company) assumes no responsibility for any occurred problem attributable to the contents of the multilingual product information.

本剤についてのお問い合わせは、お買い求めのお店又は下記にお願いいたします。

八ッ目製薬株式会社(お客様相談室)

電 話 (03) 3680-0005

受付時間 9:00~17:00 (土・日、祝日を除く)

製造販売元 ハッ目製薬株式会社

東京都江戸川区船堀2-14-14(〒134-0091)